

Erratum

The Effects of Nitric Oxide on Coagulation During Simulated Extracorporeal Membrane Oxygenation

Eric D. Rauch, BS, CCP; Alfred H. Stammers, MSA, CCP; Brian L. Mejak, BS, CCP;
See N. Vang, BS, CCP; Thomas W. Viessman, BSMT, CCP

Division of Clinical Perfusion Education, School of Allied Health Professions, University of Nebraska Medical Center, Omaha, Nebraska

[Article in The Journal of Extra-Corporeal Technology, Volume 32, Number 4, December 2000]

The correct version of each legend for Figures 2-8 of this article should read:




Figure 2. Platelet count data in mm³ displayed across the five sample times.




Figure 3. Plasma-free hemoglobin data in mg dL⁻¹ displayed across the five sample times.




Figure 4. Methemoglobin data in mg dL⁻¹ displayed across the five sample times.




Figure 5. Maximum amplitude data in mm displayed across the five sample times.




Figure 6. Platelet counts (mm³) and TEG indices displayed across the five sample times.




Figure 7. Scanning electron micrograph of the silicone membrane oxygenator at 40,000 magnification; representation of a positive finding in the control group.




Figure 8. Scanning electron micrograph of the silicone membrane oxygenator at 40,000 magnification; representation of a negative finding in the treatment group.